

MEN'S ATTITUDE SURVEY

MARKING INSTRUCTIONS

- Make dark marks that fill the circle completely.
- Make clean erasures.
- Make NO stray marks.
- DO NOT fold this form.
- Fill in one circle for each question.

INCORRECT MARKS



CORRECT MARK



| ID NUMBER | | | |
|-----------|---|---|---|
| | | | |
| 0 | 0 | 0 | 0 |
| 1 | 1 | 1 | 1 |
| 2 | 2 | 2 | 2 |
| 3 | 3 | 3 | 3 |
| 4 | 4 | 4 | 4 |
| 5 | 5 | 5 | 5 |
| 6 | 6 | 6 | 6 |
| 7 | 7 | 7 | 7 |
| 8 | 8 | 8 | 8 |
| 9 | 9 | 9 | 9 |

| VISIT NO. |
|-----------|
| 470 |
| 0 |
| 1 |
| 2 |
| 3 |
| 4 |
| 5 |
| 6 |
| 7 |
| 8 |
| 9 |

| DATE | | |
|------|-----|------|
| MO. | DAY | YEAR |
| Jan | | 07 |
| Feb | | |
| Mar | 00 | 00 |
| Apr | 01 | 01 |
| May | 02 | 02 |
| June | 03 | 03 |
| July | 04 | 04 |
| Aug | 05 | 05 |
| Sept | 06 | 06 |
| Oct | 07 | 07 |
| Nov | 08 | 08 |
| Dec | 09 | 09 |

Combination Drug Treatments for HIV

For these questions “**combination drug treatments**” refers to combinations of drugs—including protease inhibitors—used to keep people who are **already infected** with HIV from getting sick with AIDS. In addition, these drugs may also be used by an HIV negative person following a possible exposure to HIV (e.g., through unsafe sex or needle sharing) to try and **prevent** infection with HIV. This is called “**post-exposure prevention**,” or **PEP** for short. Sometimes, these drugs may be used by an HIV negative person before possible exposure to HIV to try and prevent infection with HIV. This is called “**pre-exposure prevention**,” or **PREP** for short.

- How much have you heard about using combination drug treatments to try to prevent infection after a possible exposure to HIV (PEP)?
 - 0 nothing
 - 1 a little
 - 2 some
 - 3 quite a bit
 - 4 a lot
- How many people do you know who have used combination drug treatments to try to prevent infection after a possible exposure to HIV (PEP)?
 - none
 - 1 or 2
 - 3 or more
- Have you ever used combination drug treatments to try to prevent infection after a possible exposure to HIV (PEP)?
 - no
 - yes
- How much have you heard about using combination drug treatments to try to prevent infection before possible exposure to HIV (PREP)?
 - 0 nothing
 - 1 a little
 - 2 some
 - 3 quite a bit
 - 4 a lot
- How many people do you know who have used combination drug treatments to try to prevent infection before possible exposure to HIV (PREP)?
 - none
 - 1 or 2
 - 3 or more
- Have you ever used combination drug treatments to try to prevent infection before possible exposure to HIV (PREP)?
 - no
 - yes
- A. Have you and your main partner discussed rules for sex with other people?
 - no → SKIP TO Q8
 - yes
 - no main partner → SKIP TO Q8

If yes . . . does your agreement include:
(Mark all that apply)

- B. no sex outside the relationship
- C. no oral sex with another man
- D. no anal sex with another man
- E. discussed topic, but no rules established
- F. anal sex with others, but only with condom
- G. other, please describe: _____

Personal Attitudes

8. PLEASE MARK ONE NUMBER TO SHOW HOW MUCH YOU AGREE WITH EACH STATEMENT, GIVING YOUR FIRST REACTION.

| | Do not agree at all | | | | Strongly Agree |
|--|---------------------|---|---|---|----------------|
| | ① | ② | ③ | ④ | ⑤ |
| 1. HIV is no longer the threat it used to be. | ① | ② | ③ | ④ | ⑤ |
| 2. I am somewhat more willing to take a chance of getting infected or infecting someone else now that combination drug treatments are available. | ① | ② | ③ | ④ | ⑤ |
| 3. I regard myself as a “barebacker”. | ① | ② | ③ | ④ | ⑤ |
| 4. Because of combination drug treatments for HIV, I am less concerned about becoming HIV+ or infecting someone. | ① | ② | ③ | ④ | ⑤ |
| 5. I feel tired of always having to monitor my sexual behavior. | ① | ② | ③ | ④ | ⑤ |
| 6. HIV is now a controllable disease like diabetes. | ① | ② | ③ | ④ | ⑤ |
| 7. The pleasure of skin to skin sex outweighs the threat of infection or reinfection with HIV. | ① | ② | ③ | ④ | ⑤ |
| 8. I find it difficult to maintain my commitment to safer sex. | ① | ② | ③ | ④ | ⑤ |
| 9. I like wild “uninhibited” sexual encounters. | ① | ② | ③ | ④ | ⑤ |
| 10. I like new and exciting sexual experiences and sensations. | ① | ② | ③ | ④ | ⑤ |
| 11. I am a lot less worried about sexual “slipping” now that treatments may be given after unprotected sex. | ① | ② | ③ | ④ | ⑤ |
| 12. The availability of combination drug treatments for HIV makes me less worried about having unprotected sex. | ① | ② | ③ | ④ | ⑤ |
| 13. Reinfection with HIV is a serious concern for me. | ① | ② | ③ | ④ | ⑤ |
| 14. By taking combination drug treatments, an HIV+ person decreases the chance that he/she will infect a partner with HIV. | ① | ② | ③ | ④ | ⑤ |
| 15. I am less concerned about having unprotected anal sex now that combination drug treatments are available. | ① | ② | ③ | ④ | ⑤ |
| 16. An HIV+ person whose level of virus in the blood is undetectable is unlikely to transmit HIV to a sex partner. | ① | ② | ③ | ④ | ⑤ |
| 17. I am interested in trying out new sexual experiences. | ① | ② | ③ | ④ | ⑤ |
| 18. It takes a lot of effort to keep my sexual behavior safe. | ① | ② | ③ | ④ | ⑤ |
| 19. No one has demonstrated that strains of HIV that will make someone resistant to multiple drugs can be transmitted sexually. | ① | ② | ③ | ④ | ⑤ |
| 20. I am very concerned about the transmission of Hepatitis C during unprotected sex. | ① | ② | ③ | ④ | ⑤ |