

14. PERIPHERAL NEUROPATHY SCREENING

RIGHT

a1. Perception of vibration (at great toe) NO
 YES
 Unable to evaluate
 REFUSED

IF YES: Vibration was felt for: >10 sec. (normal)
 5-10 sec. (mild loss)
 >0 and <5 sec. (moderate loss)
 0 sec. (severe loss)

RIGHT

b1. Deep tendon reflexes (ankle reflexes) NO, reflexes absent
 YES, reflexes present
 Unable to evaluate
 REFUSED

IF YES: Reflexes felt were: Hypoactive
 Normal deep tendon reflexes
 Hyperactive deep tendon reflexes (e.g., with prominent spread)
 Clonus

LEFT

a2. Perception of vibration (at great toe) NO
 YES
 Unable to evaluate
 REFUSED

IF YES: Vibration was felt for: >10 sec. (normal)
 5-10 sec. (mild loss)
 >0 and <5 sec. (moderate loss)
 0 sec. (severe loss)

LEFT

b2. Deep tendon reflexes (ankle reflexes) NO, reflexes absent
 YES, reflexes present
 Unable to evaluate
 REFUSED

IF YES: Reflexes felt were: Hypoactive
 Normal deep tendon reflexes
 Hyperactive deep tendon reflexes (e.g., with prominent spread)
 Clonus

Additional Comments:

Horizontal lines for additional comments.

PERF

5/8" Glued

PERF

LIPODYSTROPHY PHYSICAL EXAMINATION

1. Height:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

2. Mid-Arm Girth:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

(see instructions)

3. Chest Girth:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

4. Waist Girth:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

5. Hip Girth:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

6. Thigh Girth

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

LIPODYSTROPHY MEASURER CODE	
0	0
10	1
20	2
30	3
40	4
50	5
60	6
70	7
80	8
90	9

7. Thigh Skinfold:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

8. Triceps Skinfold:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

9. Subscapular Skinfold:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

10. Biceps Skinfold:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

11. Breast Skinfold:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

12. Abdominal Skinfold:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

13. Suprailiac Skinfold:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

14. Fat Wasting (see severity definitions below):

*If None or Refused, go to next question.
If Yes, indicate severity of symptom.*

	Severity*		
	Mild	Moderate	Severe
1) Facial fat loss (sunken cheeks)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2) Arms	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3) Legs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4) Buttocks	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

15. Fat Accumulation:

*If None or Refused, go to next question.
If Yes, indicate severity of symptom.*

	Severity*		
	Mild	Moderate	Severe
1) Moon facies	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2) Abdomen	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3) Back of Neck	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4) Breasts	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

16. Other physical exam findings noted related to fat distribution:

Specify:

* Definitions:

- None: Patient does not exhibit any signs of fat maldistribution. (Not noted by patient or clinician)
- Mild: Mild signs noted only after close inspection by patient or clinician.
- Moderate: Signs of fat maldistribution are noticed by patient or clinician without specifically looking for it. Patient may complain that current clothing has become tighter.
- Severe: Signs of fat maldistribution easily noted by casual observer. Symptoms have required a change in size of clothing or undergarments worn.