

LIPODYSTROPHY PHYSICAL EXAMINATION

1. Height:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

2. Mid-Arm Girth:

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

(see instructions)

3. Chest Girth:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

4. Waist Girth:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

5. Hip Girth:

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

6. Thigh Girth

0	0	0	0
1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

LIPODYSTROPHY MEASURER CODE		
0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

7. Fat Wasting (see severity definitions below):

If None or Refused, go to next question.
If Yes, indicate severity of symptom.

Severity*

Mild Moderate Severe

Refused None Yes

1) Facial fat loss (sunken cheeks)

2) Arms

3) Legs

4) Buttocks

8. Fat Accumulation:

If None or Refused, go to next question.
If Yes, indicate severity of symptom.

Severity*

Mild Moderate Severe

Refused None Yes

1) Moon facies

2) Abdomen

3) Back of Neck

4) Breasts

* Definitions:

None: Patient does not exhibit any signs of fat maldistribution. (Not noted by patient or clinician)

Mild: Mild signs noted only after close inspection by patient or clinician.

Moderate: Signs of fat maldistribution are noticed by patient or clinician without specifically looking for it. Patient may complain that current clothing has become tighter.

Severe: Signs of fat maldistribution easily noted by casual observer. Symptoms have required a change in size of clothing or undergarments worn.

Q7. Fat Wasting:

The examiner observes and grades the lipoatrophy (both facial and limb) according to the following standards:

For facial lipoatrophy:

- mild — clearly visible deepened nasolabial folds
- moderate — evidence of “hollowing out” of cheeks
- severe — hollowed cheek areas with underlying muscle clearly visible

For limb (arms and legs) lipoatrophy:

- mild — increased prominence of veins
- moderate — increased prominence of both veins and muscles
- severe — a + b with overall thinning appearance of the limb

9. Other physical exam findings noted related to fat distribution:

Specify: