

7. OROPHARYNGEAL

NO YES REFUSED

a. Consistent with oral thrush/candidiasis (ENTTH_48)

IF YES:

- KOH negative
-OR-
 KOH positive
 Not performed

ENTKO_48

b. Consistent with herpetic lesions (ENTHP_48)**c. Gingivitis/gum disease** (ENTGG_48)**d. Oral hairy leukoplakia** (ENTLE_48)**e. Other (please describe below)** (ENTOT_48)

8. EYES

NO YES REFUSED

a. Conjunctiva

1) Redness

2) Discharge

EYRED_48

EYDIS_48

b. Scleral icterus

EYSCI_48

c. Other (please describe below)

EYOTH_48

9. LYMPH NODES

NO YES REFUSED

a. Are there any nodes present
(excluding inguinal and femoral)
which are ≥ 1 cm?

LYND1_48

SKIP TO Q 10 ←

b. Presence of node ≥ 1 cm

1) Occipital

Right OCCR_48

Left OCCL_48

2) Post. auricular

Right POAUR_48

Left POAUL_48

3) Pre-auricular

Right PRAUR_48

Left PRAUL_48

4) Submental/submandibular

Right SUBMR_48

Left SUBML_48

5) Ant. cervical

Right ACERR_48

Left ACERL_48

6) Post. cervical

Right PCERR_48

Left PCERL_48

7) Supraclavicular

Right SCLVR_48

Left SCLVL_48

8) Axillary

Right AXILR_48

Left AXILL_48

9) Epitrochlear

Right EPTRR_48

Left EPTRL_48

c. What is the diameter of the largest node present?

LNODD_48

- 1-2 cm 2.1-4 cm >4 cm

d. Are any of the nodes tender?NO YES
TENND_48**e. Are any of the nodes matted?**

MATND_48

LIPODYSTROPHY PHYSICAL EXAMINATION

1. Height:

_____ cm
HEIGHTCM_48

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

2. Mid-Arm Girth:

_____ cm
LDMID_48

1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

(see instructions)

3. Chest Girth:

_____ cm
LDCHE_48

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

4. Waist Girth:

_____ cm
LDWAI_48

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

5. Hip Girth:

_____ cm
LDHIP_48

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

6. Thigh Girth:

_____ cm
LDTHI_48

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

LIPODYSTROPHY MEASURER CODE

LPEXN_48

2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

7. Fat Wasting (see severity definitions below):

If None or Refused, go to next question.
 If Yes, indicate severity of symptom.

	Refused	None	Yes	Mild	Moderate	Severe
1) Facial fat loss (sunken cheeks)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2) Arms	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3) Legs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4) Buttocks	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

8. Fat Accumulation:

If None or Refused, go to next question.
 If Yes, indicate severity of symptom.

	Refused	None	Yes	Mild	Moderate	Severe
1) Moon facies	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2) Abdomen	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3) Back of Neck	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4) Breasts	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

* Definitions:

- None:** Patient does not exhibit any signs of fat maldistribution. (Not noted by patient or clinician)
- Mild:** Mild signs noted only after close inspection by patient or clinician.
- Moderate:** Signs of fat maldistribution are noticed by patient or clinician without specifically looking for it. Patient may complain that current clothing has become tighter.
- Severe:** Signs of fat maldistribution easily noted by casual observer. Symptoms have required a change in size of clothing or undergarments worn.

Q7. Fat Wasting:

The examiner observes and grades the lipoatrophy (both facial and limb) according to the following standards:

For facial lipoatrophy:

- a) mild — clearly visible deepened nasolabial folds
- b) moderate — evidence of “hollowing out” of cheeks
- c) severe — hollowed cheek areas with underlying muscle clearly visible

For limb (arms and legs) lipoatrophy:

- a) mild — increased prominence of veins
- b) moderate — increased prominence of both veins and muscles
- c) severe — a + b with overall thinning appearance of the limb

9. Other physical exam findings noted related to fat distribution: **PCFAT_48**

Specify: