



### 7. OROPHARYNGEAL

NO YES REFUSED

a. Consistent with oral thrush/candidiasis (ENTTH\_49)

IF YES:

- KOH negative  
 -OR-  
 KOH positive  
 Not performed

ENTKO\_49

b. Consistent with herpetic lesions (ENTHP\_49)

c. Gingivitis/gum disease (ENTGG\_49)

d. Oral hairy leukoplakia (ENTLE\_49)

e. Other (please describe below) (ENTOT\_49)

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

### 8. EYES

NO YES REFUSED

a. Conjunctiva

- 1) Redness (EYRED\_49)  
 2) Discharge (EYDIS\_49)

b. Scleral icterus (EYSCI\_49)

c. Other (please describe below) (EYOTH\_49)

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

### 9. LYMPH NODES

NO YES REFUSED

a. Are there any nodes present (excluding inguinal and femoral) which are ≥1 cm? (LYND1\_49)

**SKIP TO Q 10** ←

b. Presence of node ≥1 cm

- |                            |       |          |
|----------------------------|-------|----------|
| 1) Occipital               | Right | OCCR_49  |
|                            | Left  | OCCL_49  |
| 2) Post. auricular         | Right | POAUR_49 |
|                            | Left  | POAUL_49 |
| 3) Pre-auricular           | Right | PRAUR_49 |
|                            | Left  | PRAUL_49 |
| 4) Submental/submandibular | Right | SUBMR_49 |
|                            | Left  | SUBML_49 |
| 5) Ant. cervical           | Right | ACERR_49 |
|                            | Left  | ACERL_49 |
| 6) Post. cervical          | Right | PCERR_49 |
|                            | Left  | PCERL_49 |
| 7) Supraclavicular         | Right | SCLVR_49 |
|                            | Left  | SCLVL_49 |
| 8) Axillary                | Right | AXILR_49 |
|                            | Left  | AXILL_49 |
| 9) Epitrochlear            | Right | EPTRR_49 |
|                            | Left  | EPTRL_49 |

c. What is the diameter of the largest node present? (LNODD\_49)

- 1-2 cm    2.1-4 cm    >4 cm

d. Are any of the nodes tender? (TENND\_49)

e. Are any of the nodes matted? (MATND\_49)



14. PERIPHERAL NEUROPATHY SCREENING

RIGHT

a1. Perception of vibration (at great toe)  NO  
 YES  
 Unable to evaluate  
 REFUSED  
PNPVR\_49

IF YES: Vibration was felt for: →  >10 sec. (normal)  
 5–10 sec. (mild loss)  
PNVTR\_49  >0 and <5 sec. (moderate loss)  
 0 sec. (severe loss)

LEFT

a2. Perception of vibration (at great toe)  NO  
 YES  
 Unable to evaluate  
 REFUSED  
PNPVL\_49

IF YES: Vibration was felt for: →  >10 sec. (normal)  
 5–10 sec. (mild loss)  
PNVTL\_49  >0 and <5 sec. (moderate loss)  
 0 sec. (severe loss)

RIGHT

b1. Deep tendon reflexes (ankle reflexes)  NO, reflexes absent  
 YES, reflexes present  
 Unable to evaluate  
 REFUSED  
PNTRR\_49

IF YES: Reflexes felt were: →  Hypoactive  
 Normal deep tendon reflexes  
PNTTR\_49  Hyperactive deep tendon reflexes  
(e.g., with prominent spread)  
 Clonus

LEFT

b2. Deep tendon reflexes (ankle reflexes)  NO, reflexes absent  
 YES, reflexes present  
 Unable to evaluate  
 REFUSED  
PNTRL\_49

IF YES: Reflexes felt were: →  Hypoactive  
 Normal deep tendon reflexes  
PN TTL\_49  Hyperactive deep tendon reflexes  
(e.g., with prominent spread)  
 Clonus

Additional Comments:

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



# LIPODYSTROPHY PHYSICAL EXAMINATION

## 1. Height:

HEIGHCM\_49

(1)	(1)	(1)	(1)
(2)	(2)	(2)	(2)
(3)	(3)	(3)	(3)
(4)	(4)	(4)	(4)
(5)	(5)	(5)	(5)
(6)	(6)	(6)	(6)
(7)	(7)	(7)	(7)
(8)	(8)	(8)	(8)
(9)	(9)	(9)	(9)

(see instructions)

## 2. Mid-Arm Girth:

LDMID\_49

(1)	(1)	(1)
(2)	(2)	(2)
(3)	(3)	(3)
(4)	(4)	(4)
(5)	(5)	(5)
(6)	(6)	(6)
(7)	(7)	(7)
(8)	(8)	(8)
(9)	(9)	(9)

(see instructions)

## 3. Chest Girth:

LDLCE\_49

(2)	(2)	(2)	(2)
(3)	(3)	(3)	(3)
(4)	(4)	(4)	(4)
(5)	(5)	(5)	(5)
(6)	(6)	(6)	(6)
(7)	(7)	(7)	(7)
(8)	(8)	(8)	(8)
(9)	(9)	(9)	(9)

(see instructions)

## 4. Waist Girth:

LDWAI\_49

(2)	(2)	(2)	(2)
(3)	(3)	(3)	(3)
(4)	(4)	(4)	(4)
(5)	(5)	(5)	(5)
(6)	(6)	(6)	(6)
(7)	(7)	(7)	(7)
(8)	(8)	(8)	(8)
(9)	(9)	(9)	(9)

(see instructions)

## 5. Hip Girth:

LDHIP\_49

(2)	(2)	(2)	(2)
(3)	(3)	(3)	(3)
(4)	(4)	(4)	(4)
(5)	(5)	(5)	(5)
(6)	(6)	(6)	(6)
(7)	(7)	(7)	(7)
(8)	(8)	(8)	(8)
(9)	(9)	(9)	(9)

(see instructions)

## 6. Thigh Girth

LDTHI\_49

(1)	(1)	(1)	(1)
(2)	(2)	(2)	(2)
(3)	(3)	(3)	(3)
(4)	(4)	(4)	(4)
(5)	(5)	(5)	(5)
(6)	(6)	(6)	(6)
(7)	(7)	(7)	(7)
(8)	(8)	(8)	(8)
(9)	(9)	(9)	(9)

(see instructions)

## LIPODYSTROPHY MEASURER CODE

LPEXN\_49

(1)	(1)	(1)
(2)	(2)	(2)
(3)	(3)	(3)
(4)	(4)	(4)
(5)	(5)	(5)
(6)	(6)	(6)
(7)	(7)	(7)
(8)	(8)	(8)
(9)	(9)	(9)

## 7. Fat Wasting (see severity definitions below):

If None or Refused, go to next question.  
If Yes, indicate severity of symptom.

Refused None Yes

Severity\*  
Mild Moderate Severe

1) Facial fat loss (sunken cheeks)

FWFAC\_49

SWFAC\_49

2) Arms

FWARM\_49

SWARM\_49

3) Legs

FWLEG\_49

SWLEG\_49

4) Buttocks

FWBUT\_49

SWBUT\_49

## 8. Fat Accumulation:

If None or Refused, go to next question.  
If Yes, indicate severity of symptom.

Refused None Yes

Severity\*  
Mild Moderate Severe

1) Moon facies

FAMOO\_49

SWMOO\_49

2) Abdomen

FAABD\_49

SWABD\_49

3) Back of Neck

FANCK\_49

SWNCK\_49

4) Breasts

FABRS\_49

SWBRS\_49

### \* Definitions:

- None: Patient does not exhibit any signs of fat maldistribution. (Not noted by patient or clinician)
- Mild: Mild signs noted only after close inspection by patient or clinician.
- Moderate: Signs of fat maldistribution are noticed by patient or clinician without specifically looking for it. Patient may complain that current clothing has become tighter.
- Severe: Signs of fat maldistribution easily noted by casual observer. Symptoms have required a change in size of clothing or undergarments worn.

### Q7. Fat Wasting:

The examiner observes and grades the lipoatrophy (both facial and limb) according to the following standards:

For facial lipoatrophy:

- a) mild — clearly visible deepened nasolabial folds
- b) moderate — evidence of "hollowing out" of cheeks
- c) severe — hollowed cheek areas with underlying muscle clearly visible

PCFAT\_49  
center option

For limb (arms and legs) lipoatrophy:

- a) mild — increased prominence of veins
- b) moderate — increased prominence of both veins and muscles
- c) severe — a + b with overall thinning appearance of the limb

## 9. Other physical exam findings noted related to fat distribution:

Specify: