











# LIPODYSTROPHY PHYSICAL EXAMINATION

1. Height:     cm  
**HEIGHCM\_53**

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

2. Mid-Arm Girth:     cm  
**LDMID\_53**

1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

(see instructions)

3. Chest Girth:     cm  
**LDLCH\_53**

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

4. Waist Girth:     cm  
**LDWAI\_53**

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

5. Hip Girth:     cm  
**LDHIP\_53**

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

6. Thigh Girth:     cm  
**LDTHI\_53**

1	1	1	1
2	2	2	2
3	3	3	3
4	4	4	4
5	5	5	5
6	6	6	6
7	7	7	7
8	8	8	8
9	9	9	9

(see instructions)

LIPODYSTROPHY MEASURER CODE  
**LPEXN\_53**

0	0	0
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9

7. Fat Wasting (see severity definitions below):

If None or Refused, go to next question.  
 If Yes, indicate severity of symptom.

	Refused	None	Yes	Mild	Moderate	Severe
1) Facial fat loss (sunken cheeks)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2) Arms	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3) Legs	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4) Buttocks	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

8. Fat Accumulation:

If None or Refused, go to next question.  
 If Yes, indicate severity of symptom.

	Refused	None	Yes	Mild	Moderate	Severe
1) Moon facies	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
2) Abdomen	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
3) Back of Neck	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
4) Breasts	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

\* Definitions:

- None:** Patient does not exhibit any signs of fat maldistribution. (Not noted by patient or clinician)
- Mild:** Mild signs noted only after close inspection by patient or clinician.
- Moderate:** Signs of fat maldistribution are noticed by patient or clinician without specifically looking for it. Patient may complain that current clothing has become tighter.
- Severe:** Signs of fat maldistribution easily noted by casual observer. Symptoms have required a change in size of clothing or undergarments worn.

Q7. Fat Wasting:

The examiner observes and grades the lipoatrophy (both facial and limb) according to the following standards:

For facial lipoatrophy:

- a) mild – clearly visible deepened nasolabial folds
- b) moderate – evidence of “hollowing out” of cheeks
- c) severe – hollowed cheek areas with underlying muscle clearly visible

For limb (arms and legs) lipoatrophy:

- a) mild – increased prominence of veins
- b) moderate – increased prominence of both veins and muscles
- c) severe – a + b with overall thinning appearance of the limb

9. Other physical exam findings noted related to fat distribution: **PCFAT\_53**

Specify: